



Advancing Eco Agriculture

For more information, refer to the AEA Product Usage Guide at [advancingcoag.com/product-usage-guide](http://advancingcoag.com/product-usage-guide)



## 2-2-2

### Guaranteed analysis:

Nitrogen (N).....	2%
1.5% Water Soluble Nitrogen	
0.5% Water Insoluble Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	2%
Soluble Potash (K <sub>2</sub> O).....	2%
Calcium (Ca).....	1.5%
Magnesium (Mg).....	0.5%
Sulfur (S).....	1%

**Derived from:** Corn steep powder, calcium silicate, magnesium sulfate, kelp extract (potassium hydroxide)

### Net Contents:

- 1 gallon (3.78L)       55 gallon (208.2L)
- 9.9 lbs (4.49 kg)      544.5 lbs (244.5 kg)
- 5 gallon (18.92L)     275 gallon (1041L)
- 49.5 lbs (22.4 kg)    2722.5 lbs (1234.9 kg)
- 30 gallon (113.56L)
- 297 lbs (134.7kg)

### Density:

9.9 lbs. per gallon at 68°F

F3481

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

# MacroPak™

## An all-purpose macronutrient blend

### Application Rates And Timing

**Foliar:** Apply up to 2 quarts per acre every 7 to 10 days

**Fertigation:** Apply up to 3 gallons per acre every 7 to 10 days

**Sidedress:** Apply up to 2 gallons per acre

Check with your crop consultant for specific application rates and other uses.

The product may not be applied near water, storm drains or drainage ditches, or to any impervious surface. The product may not be applied if heavy rain is expected. The product may only be applied to the intended application site.

### Handling

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

### When Tank Mixing

A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before or during application, there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

**Disclaimer:** AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



**Advancing Eco Agriculture, Inc.**

4551 Parks West Road, Middlefield, OH 44062  
[www.advancingcoag.com](http://www.advancingcoag.com) • 800-495-6603