



Advancing Eco Agriculture

For more information, refer to the AEA Product Usage Guide at advancingecoag.com/product-usage-guide



Guaranteed analysis:

Sulfur (S)	1.5%
Cobalt (Co)	0.8%
Copper (Cu)	0.1%
Manganese (Mn)	0.4%
Molybdenum (Mo)	0.5%
Zinc (Zn)	0.3%

Derived from:

Cobalt Sulfate, Copper Sulfate, Manganese Sulfate, Sodium Molybdate, Zinc Sulfate

WARNING: Contains Molybdenum. The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals

Net Contents:

- 1 gallon (3.78L) 55 gallon (208.2L)
- 9.3 lbs (3.9 kg) 511.5 lbs (219.5 kg)
- 5 gallon (18.92L) 275 gallon (1041L)
- 46.5 lbs (19.9 kg) 2557.5 lbs (1057.8 kg)
- 30 gallon (113.56L)
- 279 lbs (119.7 kg)

Density:

9.3 lbs. per gallon at 68°F

F3481

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

SeedFlare

Mineral Seed Treatment

Application Rates And Timing

Spray SeedFlare directly onto the seed without diluting it with water to avoid softening the seed. Seed can be treated well ahead of planting.

Can be applied with a commercial seed treater. Can also be applied as a spray.

- 2-8 quarts per ton of seed
- 1.6-6.4 ounces per 50lb of seed
- 1-3.75 milliliters per pound of seed

The product may not be applied near water, storm drains or drainage ditches, or to any impervious surface. The product may not be applied if heavy rain is expected. The product may only be applied to the intended application site.

Handling:

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When Tank Mixing:

A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before or during application, there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

WARNING | DANGER



- Suspected of damaging fertility or the unborn child.
- May cause damage to organs through prolonged or repeated exposure.

California Proposition 65 Warning: This product contains an ingredient (cobalt sulfate) known to the State of California to cause cancer.

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Advancing Eco Agriculture, Inc.
4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603