



Advancing Eco Agriculture, Inc.

For more information, refer to the AEA Product Usage Guide at advancingecoag.com/product-usage-guide



Guaranteed analysis:

Iron (Fe)3%

Derived from: Ferrous Sulfate Heptahydrate

Net Contents:

- 1 gallon (3.78L) 55 gallon (208.2L)
9.2 lbs (4.17 kg) 506 lbs (229.5 kg)
- 5 gallon (18.92L) 275 gallon (1041L)
46 lbs (20.8 kg) 2530 lbs (1147.5 kg)
- 30 gallon (113.56L)
276 lbs (125 kg)

Density:

9.2 lbs. per gallon at 68°F

F3481

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

Rebound™ Iron

Nutrient Solution

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules for certified organic crop production. Verify with your organic certifier before using.

Application Rates And Timing

This material is intended for soil and foliar applications and should only be used in accordance with the manufacturer's instructions. Check with your crop consultant for specific application rates.

Soil: Apply up to 1 gallon per acre.

Broad acre crop foliar: Apply up to 1 gallon per acre, every 3-4 weeks.

Fruit and vegetable foliar: Apply up to 1 gallon per acre per week.

Irrigation systems: Apply up to 1 gallon per acre every 10-14 days.

Handling

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

The product may not be applied near water, storm drains or drainage ditches, or to any impervious surface. The product may not be applied if heavy rain is expected. The product may only be applied to the intended application site.

When Tank Mixing

A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before or during application, there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Guaranteed by Advancing Eco Agriculture, Inc.
4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603