



Advancing Eco Agriculture, Inc.

For more information, refer to the AEA Product Usage Guide at advancingecoag.com/product-usage-guide



Pinion

Controls diseases caused by the fungi in the order Erysiphales

Active Ingredients:

Citric Acid	1.4%
Cinnamon Oil	0.1%
*Inert Ingredients:	98.5%
TOTAL	100.00%

*Inert Ingredients: D-glucose, vinegar, lactic acid, maltodextrin and water.

Net Contents:

- 2.5 gallon (9.46 L)
21.7 lbs (9.84 kg)
- 275 gallon (1041 L)
2387 lbs (1082.72 kg)

Density:

8.68 lbs. per gallon at 68°F

Information regarding the contents and levels of metals in this product is available on the internet at: aapfco.org/metals.html

This product qualifies for exemption from EPA registration under Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA)

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use the product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

CONTAINER HANDLING

Nonrefillable Container. Do not reuse or refill this container.

If Empty:

- Place empty container in trash or offer for recycling, if available.

If Partly Filled:

- If it is necessary to dispose of unused product, please call your local solid waste agency for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

Important: Use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines.

Application Rates and Timing:

Fruits, Vegetables and Broad Acre Crops:

Dilute with water to make a 1% solution.

As a foliar preventative, apply 2 – 3 qt. per acre ~30 days after germination. If applying to fruit trees, apply once new plant growth is observed.

As a foliar treatment, following signs of disease development, apply 1 gallon per acre.

Requires 2 applications, 10–12 days apart. As a maintenance treatment, apply 2 – 3 qt. per acre ~monthly.

Spray only the target area to avoid exposure to beneficial organisms.

WARNING: Keep out of reach of children.

WARNING



- **Causes serious eye irritation**

Precautionary Statements:

Do not handle until all safety precautions have been read and understood. Wear protective gloves/protective clothing/eye protection/face protection. Avoid breathing vapors/spray.

First Aid Information:

In case of skin contact: Get medical attention if irritation or rash develops or persists. Wash with plenty of soap and water.

In case of eye contact: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If irritation persists, get medical attention.

If swallowed: Call a poison center or doctor if you feel unwell. If vomiting occurs naturally, have victim lean forward to reduce the risk of aspiration. Do NOT induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person.

Product performance may be compromised when mixed with sulfur applications or other fertilizers which have an oxidizing effect.

The product may not be applied near water, storm drains or drainage ditches, or to any impervious surface. The product may not be applied if heavy rain is expected. The product may only be applied to the intended application site.

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Manufactured by Advancing Eco Agriculture, Inc.

4551 Parks West Road, Middlefield, OH 44062
advancingecoag.com (800) 495-6603