



# PINION™

FOR HOME AND GARDEN  
READY TO USE

**CONTROLS FUNGI THAT CAN CAUSE  
HOUSE AND GARDEN PLANT DISEASES**

**Active Ingredients:**

Citric Acid ..... 0.14%

Cinnamon Oil ..... 0.01%

**\*Inert Ingredients:** ..... 99.85%

**TOTAL** ..... 100.00%

\*Inert Ingredients: Glucose, vinegar, lactic acid,  
maltodextrin, and water

This product qualifies for exemption from EPA registration under  
Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA)

**CAUTION:**

See back label for additional Precautionary and First Aid Statements

**Keep out of reach of children**



Manufactured by Advancing Eco Agriculture, Inc.  
4551 Parks West Road, Middlefield, OH 44062  
advancingecoag.com (800) 495-6603

**Net Contents: 32 oz. (946mL)**

Read entire container label before using this product.

**FIRST AID:**

**Eye Contact:** Rinse thoroughly with plenty of water for at least 15 minutes, lifting lower and upper eyelids. Consult a physician.

**Skin Contact:** Wash with soap and water. May cause an allergic skin reaction. In the case of skin irritation or allergic reactions see a physician.

**Ingestion:** Clean mouth with water and drink plenty of water afterwards.

**STORAGE AND DISPOSAL – PESTICIDE STORAGE**

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F.

Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container.

**CONTAINER HANDLING**

Nonrefillable Container. Do not reuse or refill this container.

**If Empty:** Place empty container in trash or offer for recycling, if available.

**If Partly Filled:** If it is necessary to dispose of unused product, please call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

**APPLICATION RATES AND TIMING:**

With all treatments, spray until leaves are visibly wet.

As a foliar preventative, apply ~30 days after germination, or once new plant growth is observed.

As a foliar treatment, following signs of disease development, apply immediately. May require 2 applications, 10-12 days apart.

As a maintenance treatment, apply monthly.