



PINION™

FOR HOME AND GARDEN
READY TO USE



**CONTROLS FUNGI THAT CAN CAUSE
HOUSE AND GARDEN PLANT DISEASES**

Active Ingredients:

Citric Acid	0.14%
Cinnamon Oil	0.01%
*Inert Ingredients:	99.85%
TOTAL	100.00%


*Inert Ingredients: Glucose, vinegar, lactic acid, maltodextrin, and water

This product qualifies for exemption from EPA registration under Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA)

CAUTION:

See back label for additional Precautionary and First Aid Statements

Keep out of reach of children



Manufactured by Advancing Eco Agriculture, Inc.
4551 Parks West Road, Middlefield, OH 44062
advancingecoag.com (800) 495-6603

Net Contents: 32 oz. (946mL)



Read entire container label before using this product.

FIRST AID:

Eye Contact: Rinse thoroughly with plenty of water for at least 15 minutes, lifting lower and upper eyelids. Consult a physician.

Skin Contact: Wash with soap and water. May cause an allergic skin reaction. In the case of skin irritation or allergic reactions see a physician.

Ingestion: Clean mouth with water and drink plenty of water afterwards.

STORAGE AND DISPOSAL – PESTICIDE STORAGE:

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F.

Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container.

CONTAINER HANDLING:

Nonrefillable Container. Do not reuse or refill this container.

If Empty: Place empty container in trash or offer for recycling, if available.

If Partly Filled: If it is necessary to dispose of unused product, please call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

APPLICATION RATES AND TIMING:

With all treatments, spray until leaves are visibly wet.

As a foliar preventative, apply ~30 days after germination, or once new plant growth is observed.

As a foliar treatment, following signs of disease development, apply immediately. May require 2 applications, 10-12 days apart.

As a maintenance treatment, apply monthly.