



Advancing Eco Agriculture Inc.

For more information, refer to the AEA Product Usage Guide at
advancingecoag.com/product-usage-guide



Rebound™ Boron

Nutrient Solution

Guaranteed analysis:

Boron (B)..... 2%

Derived from: Disodium octaborate tetrahydrate

WARNING: Contains Boron. The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crop(s).

Net Contents:

- | | |
|---|---|
| ☐ 1 gallon (3.78L)
8.8 lbs (3.99 kg) | ☐ 55 gallon (208.2L)
484 lbs (219.5 kg) |
| ☐ 5 gallon (18.92L)
44 lbs (19.9 kg) | ☐ 275 gallon (1041L)
2420 lbs (1097.6 kg) |
| ☐ 30 gallon (113.56L)
264 lbs (119.7kg) | |

Density:

8.8 lbs. per gallon at 68°F

F3481

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.html>

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules for certified organic crop production. Verify with your certifier before using.

Application Rates and Timing:

This material is intended for soil and foliar applications and should only be used in accordance with the manufacturer's instructions. Check with your crop consultant for more specific application rates.

Soil: Apply up to 1 gallon per acre.

Broad acre crop foliar: Apply up to 1 gallon per acre, every 3-4 weeks.

Fruit and vegetable foliar: Apply up to 1 gallon per acre per week.

Irrigation systems: Apply up to 1 gallon per acre every 10-14 days.

The product may not be applied near water, storm drains or drainage ditches, or to any impervious surface. The product may not be applied if heavy rain is expected. The product may only be applied to the intended application site.

Handling:

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When Tank Mixing:

A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before and during application, there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

WARNING



- Suspected of damaging fertility or the unborn child.
- Causes serious eye irritation.

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Guaranteed by Advancing Eco Agriculture, Inc.

4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603